

WELNIR
Cefdinir Capsule USP 300 mg
Cefdinir for Oral Suspension USP 125 mg

Formulation:

Each hard gelatin capsule contains :

Cefdinir USP
Eq. to Anhydrous Cefdinir 300 mg
Excipients q.s.
Approved colour used in empty capsule shells

Each 5 ml of the reconstituted suspension contains:

Cefdinir USP
Eq. to Anhydrous cefdinir 125 mg
Excipients q.s.
In flavoured syrupy base
Approved colour added

Cefdinir

A third generation cephalosporin that is similar to cefixime. In vitro tests show that it is more active than cefixime against S.aureus and E.faecalis but less active against Enterobacteriaceae.

Indications: Respiratory tract infections, Otitis media, skin and tissues infections and other infections caused by susceptible micro-organisms.

Precautions: Impaired renal functions, penicillin cross sensitivity; there have been reports of redish stools in patients given cefdinir with iron supplements.

Contraindications: Sensitivity to cephalosporins.

Drug Interactions: Antacids / food 1 hour before or 2 hours after administration. Renal excretion inhibited by probenecid. Disulfirum-like effect if alcohol is taken within 72 hours after administration. Increased bleeding effect with oral anticoagulants, increased nephrotoxicity with aminoglycosides.

Side effects: GI disturbances, vaginal moniliasis and vaginitis, allergic reactions, increase prothrombin time, super-infections, hepatic dysfunctions, Stevens- Johnson syndrome, erythema multiforme, fever, CNS effects.

Dose: Usual oral dose of 600 mg daily as a single dose or in two divided doses. Twice daily dosing should be used for pneumonia and skin infections. The dose of cefdinir may need to be reduced in patients with renal impairment. Children from 6 months of age may be given 14mg/kg daily as a single dose or in two divided doses (to a maximum daily dose of 600 mg)

Keep the medicine out of reach of children.

Storage: Store below 25°C, Protected from light & moisture.

Presentation:

1x10 Alu-Alu blister in a unit carton with package insert.

Glass Bottle containing powder for preparation of 60 ml suspension.

Mfg. Lic. No. : MB/06/308

Mfg. In India by :

M/s ASSOCIATED BIOTECH

Vill krishanpura, PO Gurumajra Baddi Distt Solan.